

**TWO NEW SPECIES OF *PRATYLENCHUS* FILIPJEV, 1936 FOUND  
ASSOCIATED WITH MEDICINAL PLANTS IN UDAIPUR DISTRICT OF  
RAJASTHAN ALONG WITH  
THE REPORT OF *PRATYLENCHUS ZEA* GRAHAM,  
1951 AND *P. THORNEI* SHER AND ALLEN, 1953**

**I.L. BAIRWA\*, A.U. SIDDIQUI AND A. PARIHAR**

*Department of Nematology, Rajasthan College of Agriculture,  
Maharana Pratap University of Agriculture & Technology,  
Udaipur, Rajasthan, India*

Corresponding author's E-mail: [siddiquiau@hotmail.com](mailto:siddiquiau@hotmail.com)

**Abstract**

*Pratylenchus ispagoli* sp. n., from the rhizosphere of Ispagol (*Plantago amplexicaulis*) and *Pratylenchus turmeri* sp. n., from the rhizosphere of turmeric (*Curcuma longa*) are described and illustrated. *P. ispagoli* sp. n., is characterized by having longer stylet (17-19  $\mu$ m), anteriorly located vulva ( $V= 73.91-76.9$ ) and comparatively longer tail ( $c = 11.79-20.00$ ) whereas, *P. turmeri* sp. n. is characterized in having longer oesophagus ( $b' = 2.81-3.69$ ), anteriorly located vulva (66.67-78.71) and longer stylet (stylet = 18.0-19.5).